

STATISTICAL REPORT 2004

Heather Mitchell, Cathy Burrows, Valerie Scott

May 2005

Contact Details:

Cathy Burrows
Data Manager
VCCR
PO Box 161
Carlton South VIC 3053

Telephone: 03 9250 0371
Fax: 03 9349 1818
E-mail: cathy.burrows@vccr.org

CONTENTS

1.	Introduction	4
2.	Participation in Screening	
2.1	National Policy.....	5
2.2	By Women.....	5
2.3	Participation by Age Group	7
2.4	Participation by Division of General Practice.....	8
2.5	Participation by Region of the Department of Human Services	10
3.	Cytology Reports	
3.1	Reporting of Squamous Cells	11
3.2	Reporting of Human Papillomavirus Change	12
3.3	Reporting of Endocervical Component	12
3.4	Reporting of Other Cells (non-cervical)	12
3.5	Use of Recommendation Codes	13
4.	Histology/Colposcopy Reports	14
5.	Correlation between Cytology and Histology/Colposcopy Reports	15
6.	Frequency of Early Re-screening	16
Appendix 1	Cytology Code Schedule	17
Appendix 2	Follow-up and Reminder Schedule	18

1. INTRODUCTION

The Victorian Cervical Cytology Registry is one of eight such registries operating throughout Australia. Each State and Territory operates its own register. Victoria was the first State to establish such a register. The Victorian Registry commenced operation in late 1989 after amendments to the *Cancer (Central Registers) Act*.

The Pap Test Registries, as they are commonly known, were introduced progressively across Australia throughout the 1990s. The main reason for establishing the registries was to provide an infrastructure for an organised (or systematic) approach to cervical screening.

Specific tasks assigned to the registries were to facilitate the regular participation of women in the National Cervical Screening Program by sending reminder letters to women, and the provision of a safety-net for women with abnormal Pap smears.

Secondary functions of the registers have developed on a more regional basis. In Victoria, other work areas of the Registry include making available the known screening history of a woman to the laboratory that is reporting the current smear, the provision of quantitative data to laboratories to assist with their quality assurance programs, and the provision of aggregate data to the Commonwealth so the National Cervical Screening Program can be judged against an agreed set of performance indicators.

This Statistical Report is one in a series of annual reports that have been published since the inception of the Victorian Registry. This Statistical Report provides timely information about screening in Victoria; in most areas, the data is additional to that published by the Commonwealth. Wherever possible, the same methodology has been adopted in this Statistical Report as is used in the Commonwealth report.

Cytology registrations are complete for 2004. Most cytology reports are pre-coded by the pathology laboratory to the Registry's Cytology Code Schedule which is included as Appendix 1. The Cytology Code Schedule allows a Pap smear report to be summarised to a five digit numeric code covering the squamous cells, evidence of human papillomavirus infection, the endocervical component, other non-cervical cells, and the recommendation made by the laboratory as regards further testing. The full text of a Pap smear report is not stored at the Registry.

Histology and colposcopy registrations in this report are as notified by March 2005. A very small proportion of all histology reports made during 2004 is expected after this time. While reasonably comprehensive registration occurs for histology reports, only a minority of colposcopy results are registered, most typically when a histology report is not available. Unlike the coding of cytology reports, coding of histology and colposcopy reports is done by the staff of the Registry. As with cytology reports, the full text of the histology and colposcopy reports is not stored at the Registry.

The Reminder and Follow-up Protocol used by the Registry is shown in Appendix 2. Reminder letters are not sent to women whose Registry records indicate a past history of hysterectomy or of cervical or uterine malignancy, or to women who are over 70 years of age.

Finally, the production of this report would not be possible without the cooperation of the staff of the pathology laboratories of Victoria, the staff of the Registry, and the support of the Management Committee. Very sincere thanks are extended to the members of all these groups.

2. PARTICIPATION IN SCREENING

2.1 National Policy

Since 1991, the policy of the National Cervical Screening Program as regards the age group and time interval for screening has been as follows:

Routine screening with Pap smears should be carried out every two years for women who have no symptoms or history suggestive of cervical pathology.

All women who have ever been sexually active should commence having Pap smears between the ages of 18 to 20 years, or one to two years after first sexual intercourse, whichever is later. In some cases, it may be appropriate to start screening before 18 years of age.

Pap smears may cease at the age of 70 years for women who have had two normal Pap smears within the last five years. Women over 70 years who have never had a Pap smear, or who request a Pap smear, should be screened.¹

This policy is currently under review by the National Cervical Screening Program.

2.2 By Women

Participation in the Registry by women is voluntary. The non-participation rate in Victoria is considered to be substantially less than 1%. Where a woman objects to her Pap smear being registered, the Registry holds no information about that test.

During 2004, a total of 588,000 Pap smears were registered. This represents an increase of about 3.0% on the previous year. These 588,000 Pap smears appeared to originate from 550,000 women.

Over a three and a half week period in July/August 2004, PapScreen Victoria ran a media campaign encouraging women to have regular Pap tests. During this period, and for about eight subsequent weeks, there was a noticeable increase in the number of Pap tests registered.

The following table shows data on the number of Pap smears registered and the number of women from whom these tests appeared to originate for each year of operation of the Registry. The number of women screened in each of these years is probably a slight overestimate because of incomplete record linkage, there being no unique identifying number for each woman. Where possible, the Medicare number of women is used to assist with accurate record linkage. Since August 1999, the Registry has used SSA-Name in the matching of incoming tests to pre-existing data on the database. This has resulted in more complete record-linkage of different episodes of care for women, compared with the previous approach to record-linkage.

¹ Screening for the Prevention of Cervical Cancer. Commonwealth Department of Health and Family Services. Canberra: AGPS 1998.

Year	No. of Pap Smears Registered	No. of Women Screened
2004	588,000	550,000
2003	571,000	532,000
2002	579,000	540,000
2001	577,000	542,000
2000	572,000	532,000
1999	603,000	558,000
1998	619,000	571,000
1997	587,000	535,000
1996	616,000	560,000
1995	590,000	530,000
1994	622,000	562,000
1993	571,000	523,000
1992	542,000	497,000
1991	545,000	498,000
1990	436,000	402,000

In interpreting the information in the above table, it is important to realise that many women in Victoria are screened on an annual basis. Participation over a longer period of time than one year cannot be derived by adding the counts for individual years.

2.3 Participation by Age Group

The participation of women by age group in cervical screening is expressed as a percentage.

- The denominator is the Estimated Resident Population², after adjustment for the estimated proportion of women who have had a hysterectomy³. While the Estimated Resident Population is available on an annual basis for Victoria, information on hysterectomy fractions is collected nationally approximately every five years; specific rates for Victoria are not available.
- The numerator is estimated from the Registry database. It is the number of women resident in Victoria who had at least one Pap smear in the time period of interest and who appear to have a cervix (that is, they have not had a hysterectomy according to information held by the Registry).

It is emphasised that participation rates are necessarily imprecise and vulnerable to measurement error in both the numerator and denominator. Measurement error probably has a greater effect on the denominator than the numerator.

The biggest impact on denominator error comes from uncertainty about hysterectomy rates. The variation in hysterectomy rates between two adjacent 5 year age groups is up to 10%. The applicability of these rates (most recently collected in 2001) to current Victorian age groups is less than ideal.

Measurement error in Registry data comes from imperfect record-linkage between multiple smears from the same woman (resulting in an overestimate of the number of women screened) and from inaccuracies in the Registry database as regards whether the Pap smear was taken from a woman with or without a cervix. Only women with a cervix are considered eligible for cervical screening.

The following table shows the estimated percentage of eligible women from each decade of the target age range who had at least one Pap smear during 2004, during the two year period 2003-2004, and the three year period 2002-2004.

Age Group	% Screened in 2004	% Screened in 2003-2004	% Screened in 2002-2004
20-29	31%	54%	70%
30-39	38%	67%	81%
40-49	40%	70%	82%
50-59	41%	72%	82%
60-69	34%	60%	67%
20-69	36.7%	64.4%	77.0%

These participation rates are a little higher than those reported last year when the two and three year participation rates were 63.9% and 76.9% respectively.

² Australian Bureau of Statistics.

³ National Health Survey 2001. Australian Bureau of Statistics

2.4 Participation by Division of General Practice

The Commonwealth Department of Health and Ageing assigns almost all Victorian postcodes to a Division of General Practice. There are twenty nine Divisions of General Practice located solely within Victoria.

For this analysis, the denominator was the estimated number of eligible women resident in the postcodes of each Division in June 2003.

Division Number	Division Name	% screened 2003-2004
301	Melbourne Division of GP	65%
302	North East Valley Division of GP	70%
303	Inner Eastern Melbourne Division of GP	72%
304	Inner South East Melbourne Division of GP	70%
305	Westgate Division of GP	57%
306	Western Division of GP	62%
307	North West Melbourne Division of GP	62%
308	The Northern Division of GP, Melbourne	58%
310	Whitehorse Division of GP	67%
311	Greater South Eastern Division of GP	67%
312	Monash Division of GP	63%
313	Central Bayside Division of GP	76%
314	Knox Division of GP	64%
315	Dandenong & District Division of GP	62%
316	Mornington Peninsula Division of GP	63%
317	GP Association of Geelong	64%
318	Central Highlands Division of GP	61%
319	North-East Victorian Division of GP	67%
320	Eastern Ranges Division of GP	64%
322	South Gippsland Division of GP	62%
323	Central-West Gippsland Division of GP	63%
324	Otway Division of GP	66%
325	Ballarat & District Division of GP	60%
326	Bendigo & District Division of GP	62%
327	Goulbourn Valley Division of GP	65%
328	East Gippsland Division of GP	68%
330	Western Victorian Division of GP	61%
331	Murray Plains Division of GP	64%
332	Mallee Division of GP	64%

This type of information, being small-area data, is subject to greater measurement error than the data in Section 2.2 and 2.3. The main source of inaccuracy in the above table derives from applying national hysterectomy fractions to the relatively small female population resident in the postcodes of a Division of General Practice. For example, it is extremely unlikely that the national hysterectomy fractions are equally applicable to the postcodes of the Inner South East Melbourne Division of General Practice and the postcodes of the Mallee Division of General Practice.

Other additional (but probably lesser) sources of measurement error derive from use of the service provider's postcode of practice if the woman's residential postcode is not known to the Registry, the proportion of Victorian Pap smears reported by laboratories physically located outside of Victoria who do not report to the Registry (this will mainly effect Divisions located on the Victoria/New South Wales and Victoria/South Australia borders), and differences between the postcode assigned by the Australian Bureau of Statistics to the Estimated Resident Population data and the postcode nominated by the woman in her usual life.

For these reasons, the data in the above table should always be interpreted and used with considerable caution.

2.5 Participation by Region of the Department of Human Services

Most Victorian postcodes are assigned to a Region of the Victorian Department of Human Services. Victoria is divided into nine Regions, five in rural Victoria and four covering metropolitan Melbourne.

As in Section 2.4, the denominator for this analysis was the estimated number of eligible women resident in the postcodes of each Region in June 2003.

The problems relating to measurement error described in Section 2.4 also apply to the data in this section.

Region Name	% screened 2003-2004
Barwon South Western	64%
Eastern Metropolitan	66%
Gippsland	64%
Grampians	61%
Hume	67%
Loddon Mallee	64%
Northern Metropolitan	64%
Southern Metropolitan	67%
Western Metropolitan	60%

3. CYTOLOGY REPORTS

Cytology reports received by the Registry are coded numerically according to the following five categories of information which comprise the main aspects of a Pap smear report.

- * Squamous cell code
- * Human papillomavirus code
- * Endocervical component code
- * Other (non-cervical) cell code
- * Recommendation code

The following analyses relate only to the 484,000 Pap smears collected by general practitioners and nurses in 2004. Smears collected by obstetricians and gynaecologists and at hospital outpatient clinics have been excluded from the analyses in Section 3 as these are more likely to be reported as abnormal. These selection criteria thus approximate 'community based smears' from the general female population.

In the following tables, 'Average' refers to the frequency of use of the report codes across all Pap smears collected by general practitioners and nurses in 2004. 'Range' is the highest and lowest proportion for individual laboratories registering a minimum of 500 smears during 2004; eleven laboratories fulfilled these criteria. Five laboratories were excluded from this measurement because they reported less than 500 smears in 2004 to the Victorian Cervical Cytology Registry; three of these laboratories were either located on the border of Victoria and New South Wales or were located interstate.

3.1 Reporting of Squamous Cells

The following table shows the distribution of cytology reports for the ten squamous cell codes during 2004

Squamous Cell Code	Average	Range
Unsatisfactory	1.9%	0.8% - 3.6%
No abnormal cells	87.2%	75.9% - 92.7%
Minor reactive/inflammatory changes	4.4%	1.1% - 13.7%
Mild atypia	3.9%	1.7% - 6.8%
Inconclusive	0.5%	0.1% - 1.7%
CIN 1 (incl. equivocal and possible CIN 1)	1.4%	0.4% - 3.5%
CIN 2	0.3%	0.0% - 0.6%
CIN 3	0.3%	0.0% - 0.4%
Possible invasive cancer	0.01%	<0.00% - 0.03%
Invasive squamous cell carcinoma	0.01%	<0.00% - 0.01%

During 2000 and 2001 a decrease was observed in the proportion of smears reported as abnormal. While 5.0% of smears were reported as abnormal in 1999, this fell to 4.4% during 2000 and 2001. There was an increase to 5.3% in 2002 and a further increase to 6.4% in 2003. During 2004 the proportion remained the same at 6.4%.

The proportion of smears reported as showing definite high grade abnormality (ie CIN 2, CIN 3, possible invasive cancer, invasive squamous cell carcinoma) declined to 0.52% in 2000 and 2001, from its previous level of 0.7%. The proportion increased to 0.64% in 2002 and showed another slight increase to 0.67% in 2003. The proportion declined slightly to 0.62% in 2004.

3.2 Reporting of Human Papillomavirus Change

The following table shows the distribution of cytology reports according to cytological evidence of human papillomavirus (HPV) effect. HPV effect is sometimes referred to as 'wart virus infection'.

Human Papillomavirus Cell Code	Average	Range
HPV cell changes absent	97.5%	95.2% - 98.3%
HPV cell changes possible	0.4%	0.0% - 3.4%
HPV cell changes present	2.1%	1.1% - 3.1%

3.3 Reporting of Endocervical Component

The following table shows the distribution of cytology reports for the codes relating to the endocervical component. Smears which are known to have been collected post-hysterectomy are excluded.

Endocervical Component Code	Average	Range
No endocervical component present	20.3%	16.0% - 33.2%
Normal endocervical component	78.6%	66.7% - 82.1%
Minor reactive/inflammatory changes	0.9%	0.0% - 2.1%
Inconclusive	0.2%	0.0% - 0.3%
Mild/moderate dysplasia	<0.1%	0.0% - <0.1%
Adenocarcinoma in situ	<0.1%	0.0% - <0.1%
Possible invasive cancer	<0.1%	0.0% - <0.1%
Invasive adenocarcinoma	<0.1%	0.0% - <0.1%

In 1990, 27% of Pap smears lacked an endocervical component. This proportion gradually reduced over the next five years. Between 1995 and 1999, it remained stable at 15%. Since 2000 there has been a gradual increase from 17.3% in 2000 to 20.3% during 2004.

The reason for the increasing proportion of smears without an endocervical component is unclear.

3.4 Reporting of Other Cells (non-cervical)

99.9% of the cytology reports for smears collected by general practitioners and nurses indicated no other (non-cervical) abnormal cells were present.

Among the smears collected by general practitioners and nurses during 2004, there were 275 reports of benign change in non-cervical cells, 45 reports of malignant cells from the uterus, one report of malignant cells from the ovary and five reports of other malignant cells (such as metastatic malignancy).

3.5 Use of Recommendation Codes

Not all cytology reports include a recommendation by the laboratory about the next stage of care for the woman. 19,800 (4.1%) cytology reports issued during 2004 to general practitioners and nurses did not include a recommendation; this is a decrease from 20,100 in 2003.

In the following analysis, the statistics listed under 'Average' use data relating to Pap smears with a recommendation from all laboratories; the statistics listed under 'Range' are confined to the eleven laboratories that attached recommendations to more than 80% of their general practitioner/nurse Pap smears and where a minimum of 500 such reports were made. In calculating these percentages, the number of tests with recommendations was used as the denominator.

Recommendation Code	Average	Range
Repeat smear in 3 years	0.0%	0.0% - 0.1%
Repeat smear in 2 years	82.8%	68.6% - 85.6%
Repeat smear in 1 year	9.2%	7.2% - 18.9%
Repeat smear in 6 months	3.0%	1.3% - 7.2%
Repeat smear in 3 months	0.4%	0.0% - 3.4%
Repeat smear within 4 to 6 weeks	1.8%	0.1% - 2.6%
Referral for specialist opinion	2.1%	1.1% - 3.5%
Other	0.7%	0.0% - 5.3%

Among smears receiving a recommendation, the proportion recommending a repeat smear in two years has remained stable for the last three years. In earlier years there was a steady increase. In 1998, the figure was 72.9%, in 1999 it was 76.5%, in 2000 it was 79.1% and in 2001 it was 83.4%.

4. HISTOLOGY/COLPOSCOPY REPORTS

This section describes the histology/colposcopy reports taken during 2004 as known to the Registry. Less than one percent of these notifications were colposcopy reports.

From a total of 17,130 such reports known to the Registry, there were 14,775 where the report related to the cervix, with each woman being counted only once on the basis of her most severe report for the year. In ascertaining the most severe report for each woman, histology results took precedence over colposcopy results.

The following table shows the distribution of the further investigations for 2004.

Histology/colposcopy findings	Number	%
Invasive cancer	59	0.4%
Microinvasive cancer	43	0.3%
CIN 3 with questionable microinvasion	14	0.1%
CIN 3	1492	10.1%
CIN 2/3	271	1.8%
CIN 2	1482	10.0%
High grade - not otherwise defined	117	0.8%
CIN - not otherwise defined	16	0.1%
CIN 1	1758	11.9%
HPV effect	950	6.4%
Low grade - not otherwise defined	699	4.7%
Benign changes	5853	39.6%
Normal	1931	13.1%
Unsatisfactory	90	0.6%
TOTAL	14775	100%

Among the 59 women whose further investigations resulted in a diagnosis of invasive cervical cancer, 28 were of squamous type, 16 were adenocarcinomas, three were adenosquamous carcinomas, and 12 were other types.

Among the 43 women with microinvasive carcinoma, 35 (81%) were of squamous type and eight of adeno type.

Among the 1,492 cases of CIN 3, 1,418 (95%) were of squamous type, 44 were of adeno type and 30 were of adenosquamous type.

5. CORRELATION BETWEEN CYTOLOGY AND HISTOLOGY/COLPOSCOPY REPORTS

The following table shows the correlation between the histology/colposcopy findings and the prediction made on cytology immediately prior to the histology/colposcopy report.

The correlation is restricted to cases where the cytology was reported as abnormal in a six month period preceding the histology/colposcopy report. In cases where the histology or colposcopy report followed a negative cytology report, up to 24 months has been allowed between the cytology and the histology/colposcopy.

In interpreting this information, it is important to remember that only a minority of low grade cytology (atypia, HPV and CIN 1) is further investigated by colposcopy or biopsy, and an even smaller percentage of negative cytology reports are followed by a colposcopy or biopsy. Women who have a biopsy are likely to be an atypical subset of the whole group of women with negative or low grade cytology reports.

Histology/colposcopy findings	Cytology Prediction						
	Negative (n=6471)	Atypia (n=1368)	HPV (n=495)	CIN 1 (n=2237)	Inconcl. (n=937)	CIN 2 (n=1506)	CIN 3 (n=1371)
Cancer - invasive squamous	0.0%	0.0%	0.0%	0.0%	0.2%	0.1%	0.7%
Cancer - invasive other	0.1%	0.1%	0.0%	0.0%	0.4%	0.0%	0.1%
Cancer - microinvasive	0.0%	0.1%	0.0%	0.0%	0.2%	0.1%	1.7%
CIN 3 with questionable microinvasion	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%
CIN 3	0.6%	2.8%	3.0%	6.8%	20.2%	17.3%	54.4%
CIN 2/3	0.2%	0.8%	0.4%	1.3%	3.8%	5.4%	7.1%
CIN 2	1.1%	5.9%	7.7%	14.8%	15.9%	38.3%	16.3%
High grade- not otherwise defined	0.2%	0.7%	1.0%	1.0%	2.1%	1.6%	1.8%
CIN 1	3.7%	17.3%	21.0%	34.4%	10.5%	14.4%	5.4%
HPV effect	3.7%	11.7%	25.5%	11.5%	5.0%	5.1%	2.3%
Low grade- not otherwise defined	3.2%	10.5%	11.5%	7.1%	6.1%	3.5%	1.3%
Normal, benign	87.3%	50.1%	29.9%	22.9%	35.5%	14.2%	8.2%
TOTAL	100%	100%	100%	100%	100%	100%	100%

Notable points in the above table include the following.

The positive predictive value of a cytology report of CIN 3 for high grade histology (CIN 2, CIN 2/3, CIN 3, cancer, high grade - not otherwise defined) was 82.7% (1,134/1,371). For a cytology report of CIN 2 it was 62.8% (946/1506). For an inconclusive cytology report, it was 42.9% (402/937). These figures are similar to last year's.

6. FREQUENCY OF EARLY RE-SCREENING

While the Australian screening policy is for repeated testing every two years after a negative Pap smear report, many women are screened more frequently than this. While a small level of early re-screening can be justified on the basis of a past history of abnormality, the levels within Victoria and Australia are far in excess of this. The evidence is that early re-screening does not just occur in the months immediately prior to the two year anniversary, but rather is a steady continuum throughout the two year period after a negative Pap smear report.

In late 2000, the National Cervical Screening Program adopted the following definition of early re-screening:

Early re-screening is the repeating of a Pap smear within 21 months of a negative Pap smear report, except for women who are being followed up in accordance with the NHMRC guidelines for the management of cervical abnormalities.

This definition recognises that some re-screening may occur opportunistically between 21 and 24 months after a negative Pap smear report and this may be cost-effective.

The following table shows the number of further testings over a 21 month period for women who received a negative Pap smear report in the February of each year. The data shows that 73% of women aged 20-69 years who were screened in February 2003 had no further tests within the next 21 months. 27% of women aged 20-69 years who were screened in February 2003 underwent early re-screening.

Number of further Pap smears	2003	2002	2001	2000	1999	1998	1997	1996
No further tests	73%	69%	68%	65%	66%	63%	59%	57%
1	23%	26%	27%	29%	28%	31%	34%	34%
2	3%	3%	4%	4%	4%	5%	5%	6%
3	1%	1%	1%	1%	1%	1%	1%	2%
4	<1%	<1%	<1%	<1%	<1%	<1%	<1%	<1%
5 or more	<1%	<1%	<1%	<1%	<1%	<1%	<1%	<1%

The data in the above table shows a substantial improvement in early re-screening between 1996 and 2003. Among women screened in 1996, 43% had at least one additional Pap smear within 21 months. By 2003, this figure had fallen to 27%.

Some variation in early re-screening occurs by age group. The following table shows the proportion of women, by age group, who had early re-screening after a negative Pap smear report in February 2003.

Age Group	% with early re-screening
20-29 yrs	26%
30-39 yrs	30%
40-49 yrs	28%
50-59 yrs	27%
60-69 yrs	22%
20-69 yrs	27%

Early re-screening peaks in the age group 30-39 years and is least evident in the age group 60-69 years.

Appendix 1. Cytology Code Schedule

Squamous Cell Code	Wart Virus Code	Endocervical Cell Code	Other (non-cervical)	Recommendation Code
0. Unsatisfactory	- Not reported	- Not reported	- Not reported	- Not reported
		0. No endocervical component present		0. No recommendation
1. No abnormal cells	1. Absent	1. Normal endocervical component present	1. No other abnormal cells	1. Repeat smear - 3 yrs
2. Minor reactive and inflammatory changes	2. Possible	2. Minor reactive and inflammatory changes in endocervical component	2. Abnormal cells present - other (eg IUCD cells, endometrial hyperplasia)	2. Repeat smear - 2 yrs
3. Mild atypia	3. Present	3. Cell changes raising the possibility of a high grade lesion but specific diagnosis not possible (ie inconclusive)	3. Malignant cells present - uterine body	3. Repeat smear - 1 yr
4. Cell changes raising the possibility of a high grade lesion but specific diagnosis not possible (ie inconclusive)		4. Abnormal endocervical cells suggesting mild/moderate dysplasia	4. Malignant cells present - ovary	4. Repeat smear - 6 mths
5. Mild dysplasia (CIN 1) including equivocal or possible mild dysplasia		5. Abnormal endocervical cells suggesting severe dysplasia or adenocarcinoma in situ	5. Malignant cells present - vagina	5. Repeat smear - 3 mths
6. Moderate dysplasia (CIN 2)		6. Suspicious of invasive adenocarcinoma of endocervix (ie at least adenocarcinoma in situ with the possibility of invasion)	6. Malignant cells present - other (includes metastatic malignancy)	6. Repeat smear - 4 wks or less (including immediate repeat with or without treatment for infection, atrophy etc)
7. Severe dysplasia /carcinoma in situ (CIN 3)		7. Adenocarcinoma - invasive		7. Referral for specialist opinion (eg colposcopy)
8. Suspicious of micro-invasion or invasion (ie at least CIN 3 but possible invasion)				8. Other
9. Squamous cell carcinoma - invasive		9. Not applicable because of hysterectomy		

Appendix 2. Reminder and Follow-up Protocol

Cytology Report	Time*	Action by Victorian Cervical Cytology Registry
CIN 2, CIN 3, Cancer	3 months	Contact laboratory
	4 months	Questionnaire to practitioner Questionnaire to specialist if referred
	6 months	Telephone call to practitioner Letter to woman
	12 months	Reminder to woman
HPV effect, CIN 1, Inconclusive	9 months	Courtesy list to practitioner
	10 months	Reminder to woman
Atypia	13 months	Courtesy list to practitioner
	14 months	Reminder to woman
Negative with past history of histologically confirmed CIN 2 or 3	15 months	Reminder to woman
Negative with no past history of histologically confirmed CIN 2 or 3	27 months	Reminder to woman
Unsatisfactory	3 months	Repeat smear reminder to practitioner
	4 months	Reminder to woman

* Time intervals are determined from the date of the cytology